


重组人血小板生成素(TPO)

Thrombopoietin(TPO),Human,Recombinant

Cat. No.: MA0618 Size: 10μg


Source:	Human Cells
Description:	Recombinant Human Thrombopoietin is produced by our Mammalian expression system and the target gene encoding Ser22-Gly353 is expressed with a 6His tag at the N-terminus, 6His tag at the C-terminus.
Accession:	P40225
Known As:	Thrombopoietin; C-mpl ligand; Megakaryocyte colony-stimulating factor; Megakaryocyte growth and development factor; Myeloproliferative leukemia virus oncogene ligand; THPO
Predicted Mol Mass:	37.3 KDa
Apparent Mol Mass:	70-90 KDa, reducing conditions
Endotoxin:	< 0.01 EU/μg as determined by LAL test.
Formulation:	Lyophilized from a 0.2 μm filtered solution of 20mM Tris, 150mM NaCl, pH 8.0.
Reconstitution:	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Shipping:	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Storage:	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Background:	Thrombopoietin (TPO) is a glycoprotein hormone which belongs to the EPO/TPO family. It produced by the liver and kidney which regulates the production of platelets. TPO stimulates the production and differentiation of megakaryocytes, the bone marrow cells that bud off large numbers of platelets. Lineage-specific cytokine affects the proliferation and maturation of megakaryocytes from their committed progenitor cells. It acts at a late stage of megakaryocyte development. It may be the major physiological regulator of circulating platelets.
Purity-SDS-PAGE:	





Greater than 95% as determined by reducing SDS-PAGE.

Bioactivity-Cell Based Assay:



Measured in a cell proliferation assay using MO7E human megakaryocytic leukemic cells. The ED50 for this effect is 0.55 ng/ml.

