


重组大鼠白介素2(IL-2)

IL-2, Recombinant Rat Interleukin-2(C-6His)

Cat. No.: MA1356-1 Size: 10μg

Source:	Human Cells
Description:	Recombinant Rat Interleukin-2 is produced by our Mammalian expression system and the target gene encoding Ala21-Gln155 is expressed with a 6His tag at the C-terminus.
Accession:	P17108
Known As:	Interleukin-2; IL-2; T-cell growth factor; TCGF; Aldesleukin; IL2
Predicted Mol Mass:	16.3 KDa
Apparent Mol Mass:	14-16 KDa, reducing conditions
Endotoxin:	< 1 EU/μg as determined by LAL test.
Formulation:	Supplied as a 0.2 μm filtered solution of 5mM NaH ₂ PO ₄ , 5mM Citric acid, 150mM NaCl, pH 4.0.
Shipping:	The product is shipped on dry ice/polar packs. Upon receipt, store it immediately at the temperature listed below.
Storage:	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Background:	Interleukin-2(IL-2) is a O-glycosylated four α-helix bundle cytokine that has potent stimulatory activity for antigenactivated T cells. It is expressed by CD4+ and CD8+ T cells, γδ T cells, B cells, dendritic cells, and eosinophils. Mature rat IL-2 shares 66% and 73% amino acid sequence identity with human and mouse IL-2, respectively. The receptor for IL-2 consists of three subunits that are present on the cell surface in varying preformed complexes. IL-2 is a powerful immunoregulatory lymphokine produced by T-cells in response to antigenic or mitogenic stimulation. IL-2/IL-2R signaling is required for T-cell proliferation and other fundamental functions that are essential for the immune response. IL-2 stimulates growth and differentiation of B-cells, NK cells, lymphokine-activated killer cells, monocytes, macrophages and oligodendrocytes.
Purity-SDS-PAGE:	





Greater than 95% as determined by reducing SDS-PAGE.

S250101

大连美仑生物技术有限公司

官网: <https://www.meilunbio.com/>

电话/邮箱: 0411-62910999 sales@meilune.com

本产品仅供科研使用

