

重组大鼠肿瘤坏死因子 (TNF-)


TNF- ,Tumor Necrosis Factor Alpha,Rat,Recombinant

Cat. No.: MA1365-1 Size: 10μg

Source:	E.coli
Description:	Recombinant Rat Tumor Necrosis Factor is produced by our E.coli expression system and the target gene encoding Leu80-Leu235 is expressed.
Accession:	P16599
Known As:	Tumor Necrosis Factor; Cachectin; TNF-Alpha; Tumor Necrosis Factor Ligand Superfamily Member 2; TNF-a; Tumor Necrosis Factor; Membrane Form; Tumor Necrosis Factor; Soluble Form; Tnf; Tnfa; Tnfsf2
Predicted Mol Mass:	17.4 KDa
Apparent Mol Mass:	14 KDa, reducing conditions
Endotoxin:	< 1 EU/μg as determined by LAL test.
Formulation:	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Reconstitution:	Always centrifuge tubes before opening.Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Shipping:	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Storage:	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Background:	Tumor necrosis factor alpha (TNF-alpha, TNFSF2) is the prototypic ligand of the TNF superfamily. Rat TNF-alpha consists of a 35 amino acid (aa) cytoplasmic domain, a 21 aa transmembrane segment, and a 179 aa extracellular domain (ECD). Within the ECD, rat TNF-alpha shares 94% aa sequence identity with mouse. TNF-alpha is produced by a wide variety of immune, epithelial, endothelial, and tumor cells. TNF exists as a homotrimer and interacts with SPPL2B. TNF is mainly secreted by macrophages and can induce cell death of certain tumor cell lines. TNF is a key cytokine in the development of several inflammatory disorders. It contributes to the development of type 2 diabetes through its effects on insulin resistance and fatty acid metabolism.




Purity-SDS-PAGE:



Greater than 95% as determined by reducing SDS-PAGE. (QC verified)

Purity-SEC-HPLC:



Greater than 90% as determined by SEC-HPLC. (QC verified)

