

重组人白介素37(IL-37)


IL-37,Human;Recombinant Human Interleukin 37

Cat. No.: MA1419-1 Size: 10µg

Source:	E.coli
Description:	Recombinant Human Interleukin-37 is produced by our E.coli expression system and the target gene encoding Lys53-Asp218 is expressed.
Accession:	Q9NZH6
Known As:	Interleukin-37; FIL1 Zeta; IL-1X; Interleukin-1 Family Member 7; IL-1F7; Interleukin-1 Homolog 4; IL-1H; IL-1H4; Interleukin-1 Zeta; IL-1 Zeta; Interleukin-1-Related Protein; IL-1RP1; Interleukin-23; IL-37; IL37; FIL1Z; IL1F7; IL1H4; IL1RP1
Predicted Mol Mass:	18.7 KDa
Apparent Mol Mass:	18 KDa, reducing conditions
Endotoxin:	< 1 EU/µg as determined by LAL test.
Formulation:	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, 2mM DTT, pH 7.4.
Reconstitution:	Always centrifuge tubes before opening.Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Shipping:	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Storage:	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Background:	Human Interleukin family 1 Member 7 (IL1F7) is a member of the Interleukin 1 cytokine family. Five alternatively spliced transcript variants encoding distinct isoforms have been reported with distinct expression profiles. The longest IL1F7 transcript, referred to as IL1F7b or IL1F7 isoform 1, encodes a 218 amino acid residues proprotein containing a 45 amino acid propeptide, which is cleaved to generate mature protein. IL1F7b binds to IL18 Rα with low affinity but does not exert any IL18 agonistic or antagonistic effects. IL1F7b also binds interleukin 18 binding protein (IL-18BP), an inhibitory binding protein of interleukin 18 (IL-18), and subsequently forms a complex with IL18 receptor beta subunit, and through which it inhibits the activity of IL-18.



Purity-SDS-PAGE:



Greater than 95% as determined by reducing SDS-PAGE. (QC verified)

S250101

大连美仑生物技术有限公司

官网: <https://www.meilunbio.com/>

电话/邮箱: 0411-62910999 sales@meilune.com

本产品仅供科研使用

