

重组人成纤维细胞因子FGF-21

Fibroblast growth factor 21(FGF-21),Human,Recombinant(C-6His)


Cat. No.: MA1434-1 Size: 10µg

Source:	E.coli
Description:	Recombinant Human Fibroblast Growth Factor 21 is produced by our E.coli expression system and the target gene encoding His29-Ser209 is expressed with a 6His tag at the N-terminus.
Accession:	AAH18404.1
Known As:	Fibroblast Growth Factor 21; FGF-21; FGF21
Predicted Mol Mass:	21.69 KDa
Apparent Mol Mass:	23 KDa, reducing conditions
Endotoxin:	< 1 EU/µg as determined by LAL test.
Formulation:	Lyophilized from a 0.2 µm filtered solution of 20mM Tris-HCl, 100mM NaCl, 2mM EDTA, pH 9.0 .
Reconstitution:	Always centrifuge tubes before opening.Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Shipping:	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Storage:	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Background:	Fibroblast Growth Factor 21 (FGF21) is a growth factor that belongs to the FGF family. FGF family proteins play a central role during prenatal development and postnatal growth and regeneration of many tissues, by promoting cellular proliferation and differentiation. FGF21 is a potent activator of glucose uptake on adipocytes, protects animal from diet-induced obesity when overexpression in transgenic mice, and lower blood glucose and triglyceride levels when therapeutically administered to diabetic rodents. FGF21 is produced by hepatocytes in response to free fatty acid stimulation of a PPARα/RXR dimeric complex. This situation occurs clinically during starvation, or following the ingestion of a highly-fat/low-carbohydrate diet.Upon FGF21 secretion, white adipose tissue is induced to release free fatty acids from triglyceride stores.



Once free fatty acid reach hepatocytes, they are oxidized and reduced to acetyl-CoA. The acetyl-CoA is recombined into 4-carbon ketone bodies, release, and transported to peripheral tissue for TCA processing and energy generation.

Purity-SDS-PAGE:



Greater than 95% as determined by reducing SDS-PAGE. (QC verified)

