

重组小鼠胰岛素样生长因子结合蛋白6(IGFBP-6)


Insulin-Like Growth Factor-Binding Protein 6, Mouse, Recombinant (C-6His)

Cat. No.: MA1438-1 Size: 10 μ g

Source:	Human Cells
Description:	Recombinant Mouse Insulin-Like Growth Factor-Binding Protein 6 is produced by our Mammalian expression system and the target gene encoding Ala26-Gly238 is expressed with a 6His tag at the C-terminus.
Accession:	P47880
Known As:	Insulin-like growth factor-binding protein 6; IBP-6; IGF-binding protein 6; IGFBP-6; Igfbp6; IBP6; IGF binding protein 6; insulin-like growth factor-binding protein 6
Predicted Mol Mass:	23.7 KDa
Apparent Mol Mass:	28 KDa, reducing conditions
Endotoxin:	< 1 EU/ μ g as determined by LAL test.
Formulation:	Lyophilized from a 0.2 μ m filtered solution of PBS, pH 7.4.
Reconstitution:	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
Shipping:	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Storage:	Lyophilized protein should be stored at $\leq -20^{\circ}\text{C}$, stable for one year after receipt. Reconstituted protein solution can be stored at $2-8^{\circ}\text{C}$ for 2-7 days. Aliquots of reconstituted samples are stable at $\leq -20^{\circ}\text{C}$ for 3 months.
Background:	Insulin-like growth factors (IGFs) comprise a family of endocrine, paracrine and autocrine polypeptides consisting of the ligands IGF1 and IGF2, two receptors (IGF1R, IGF2R), at least 6 IGF-binding proteins (IGFBPs) and IGFBP proteases. Among the binding proteins, IGFBP6 is unique because of its N-terminal disulfide linkages and its marked binding preference for IGF2. It is a potent inhibitor of the interaction between IGF2 and its receptor IGF1R, thus preventing major functions of IGF2, such as induction of proliferation, differentiation, cell adhesion, or colony formation. In particular, IGFBP-6 inhibited the growth of neuroblastoma and rhabdomyosarcoma xenografts. IGFBP-6 is expressed in many tissues, including lung, liver, gut and the central nervous system.



Purity-SDS-PAGE:



Greater than 95% as determined by reducing SDS-PAGE.

