

## 重组小鼠OPG蛋白


TNFRSF11B/Osteoprotegerin/OPG, Mouse, Recombinant

Cat. No.: MA1470-1    Size: 10 $\mu$ g

<b>Source:</b>	Human Cells
<b>Description:</b>	Recombinant Mouse Tumor Necrosis Factor Receptor Superfamily Member 11B is produced by our Mammalian expression system and the target gene encoding Glu22-Leu401 is expressed with a 10His tag at the C-terminus.
<b>Accession:</b>	<a href="#">O08712</a>
<b>Known As:</b>	Tumor necrosis factor receptor superfamily member 11B; Osteoclastogenesis inhibitory factor; Osteoprotegerin; Tnfrsf11b; Ocif; Opg
<b>Predicted Mol Mass:</b>	45.1 KDa
<b>Apparent Mol Mass:</b>	50-65 KDa, reducing conditions
<b>Endotoxin:</b>	< 1 EU/ $\mu$ g as determined by LAL test.
<b>Formulation:</b>	Lyophilized from a 0.2 $\mu$ m filtered solution of PBS, pH 7.4.
<b>Reconstitution:</b>	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 $\mu$ g/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
<b>Shipping:</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
<b>Storage:</b>	Lyophilized protein should be stored at $\leq -20^{\circ}\text{C}$ , stable for one year after receipt. Reconstituted protein solution can be stored at 2-8 $^{\circ}\text{C}$ for 2-7 days. Aliquots of reconstituted samples are stable at $\leq -20^{\circ}\text{C}$ for 3 months.
<b>Background:</b>	Osteoprotegerin (OPG, Tnfrsf11b) is a secreted protein that regulates bone density. OPG is widely expressed and constitutively released as a homodimer by mesenchymal stem cells, fibroblasts and endothelial cells. Regulation of its expression by estrogen, parathyroid hormone and cytokines is complex and changes with age. OPG acts as decoy receptor for TNFSF11/RANKL and thereby neutralizes its function in osteoclastogenesis. TRAIL decreases the release of OPG from cells that express it, while OPG inhibits TRAIL-induced apoptosis. Expression of RANK L on the cell surface, and thus its ability to stimulate osteoclastogenesis, is regulated by OPG by intracellular and extracellular mechanisms. Bone homeostasis seems to depend on the local ratio between TNFSF11 and TNFRSF11B. It may also play a role in preventing arterial calcification.



**Purity-SDS-PAGE:**



Greater than 95% as determined by reducing SDS-PAGE.

