

Recombinant Human AITR Ligand/TNFSF18

Cat.No.:MGC133 size:5µg/50µg/500µg

Characteristics:

Source	<i>E.coli</i>
Species	Human
Tag	Tag free
Description	Human AITR Ligand/TNFSF18 (Glu74-Ser199)
Accession#	Q9UNG2
Predicted molecular mass	~14.3 kDa. It is composed of one non-glycosylated polypeptide chain containing 126 amino acids.

Specification:


Appearance	White powder, Colorless clear liquid after reconstitution
Purity	≥95%,by SDS-PAGE(under reducing(R)&Non-reducing conditions, visualized by Coomassie staining)
Endotoxin	< 0.01 EU/µg
Activity	The biological activity as determined by stimulation of IL-8 production using human PBMC is in a concentration range of 0.1-10 ng/ml.
Formulation	0.2 µm-filtered solution containing 20 mM PB, with 150 mM NaCl, pH 7.4.

Handling and Storage:

Reconstitution	It is recommended to redissolve in sterile deionized water. 细胞因子溶解、稀释、保存可使用即用型专用溶剂套装—PWL240（1mL无菌水+10mL 稀释液）
Shipping	Wet ice (seasonal)
Storage & Stability	36 months at -20°C to -80°C in lyophilized state 6 months at -20°C to -80°C under sterile conditions after reconstitution 7-10 days at 2°C to 8°C under sterile conditions after reconstitution Use a manual defrost freezer and avoid repeated freeze-thaw cycles.


Data:

SDS-PAGE



Recombinant Human AITR Ligand/TNFSF18 (Cat. No. MGC133) SDS-PAGE under reducing (R) & Non-reducing conditions.

Bioactivity



The biological activity as determined by stimulation of IL-8 production using human PBMC is in a concentration range of 0.1-10 ng/ml.

S260701